



The supersolid phase of matter

Giovanni Modugno

Dipartimento di Fisica e Astronomia and LENS, Università di Firenze,
and CNR-INO, sezione di Pisa

106° Congresso SIF, Sezione Giovani, 17-9-2020




*Quantum Simulations of
Insulators and Conductors*



INO-CNR
ISTITUTO
NAZIONALE DI
OTTICA



What is a supersolid?



Crystals: particles are localized in lattice sites



stiffness (nonzero rigidity) and
plasticity (nonzero stress to produce permanent
deformations)




Superfluids: particles are undistinguishable and
uniformly delocalized



supercurrents (zero viscosity) and
absence of stiffness (zero rigidity)

Supersolids: coexistence of the two states
above, due to interaction effects (no external
lattice).





Phys. Rev. 106, 161 (1957)

Unified Theory of Interacting Bosons

EUGENE P. GROSS

Brandeis University, Waltham, Massachusetts

(Received January 25, 1957)

RECENT work has contributed to the understanding of properties of helium II. Yet there is room for a unified theoretical approach to the problem of interacting bosons for both solid and liquid states.

Microscopic Theory of Superconductivity*

J. BARDEEN, L. N. COOPER, AND J. R. SCHRIEFFER

Department of Physics, University of Illinois, Urbana, Illinois

(Received February 18, 1957)

SINCE the discovery of the isotope effect, it has been known that superconductivity arises from the interaction between electrons and lattice vibrations, but it has proved difficult to construct an adequate theory based on this concept. As has been shown by

The Hamiltonian of identical bosons with the two body interaction potential $V(x)$... governs the motion of a classical wave field.

There is always a solution of uniform density

$$\psi(x) = \left(\frac{N}{L^3}\right)^{1/2} \text{ with energy } E = \frac{N}{L^3} \int V(x) d^3x \dots$$

but if $V(x)$ is negative in some region of space, there may be other solutions, such as periodic solutions with lower E than for the uniform solution.

Two sound sound modes



SOVIET PHYSICS JETP

VOLUME 29, NUMBER 6

DECEMBER 1969

QUANTUM THEORY OF DEFECTS IN CRYSTALS

A. F. ANDREEV and I. M. LIFSHITZ

Institute of Physical Problems, U.S.S.R. Academy of Sciences


Submitted January 15, 1969

Zh. Eksp. Teor. Fiz. 56, 2057–2068 (June, 1969)

At sufficiently low temperatures localized defects or impurities change into excitations that move practically freely through a crystal. As a result instead of the ordinary diffusion of defects, there arises a flow of a liquid consisting of “defectons” and “impuritons.” It is shown that at absolute zero in crystals with a large amplitude of the zero-point oscillations (for example, in crystals of the solid helium type) zero-point defectons may exist, as a result of which the number of sites of an ideal crystal lattice may not coincide with the number of atoms. The thermodynamic and acoustic properties of crystals containing zero-point defectons are discussed. Such a crystal is neither a solid nor a liquid. Two kinds of motion are possible in it; one possesses the properties of motion in an elastic solid, the second possesses the properties of motion in a liquid. Under certain conditions the “liquid” type of crystal motion possesses the property of superfluidity. Similar effects should also be observed in quasiequilibrium states containing a given number of defectons.

Two kinds of motion are possible in it; one possesses the properties of motion in an elastic solid, the second possesses the properties of motion in a liquid.

... we obtain an equation for the **acoustic vibrations of a crystal** ... and **oscillations of the crystal density** with fixed lattice sites...



Can a supersolid pass through a capillar without friction?

Non classical rotational inertia



Nobel 2003



Can a Solid Be “Superfluid”?

A. J. Leggett

School of Mathematical and Physical Sciences, University of Sussex, Falmer, Brighton, Sussex, England


(Received 15 September 1970)

It is suggested that the property of nonclassical rotational inertia possessed by superfluid liquid helium may be shared by some solids. In particular, nonclassical rotational inertia very probably occurs if the solid is Bose-condensed as recently proposed by Chester. Anomalous macroscopic effects are then predicted. However, the associated superfluid fraction is shown to be very small (probably $\lesssim 10^{-4}$) even at $T=0$, so that these effects could well have been missed. Direct tests are proposed.

Superfluids: macroscopic wavefunction

$$\Psi_0(r) = |\Psi_0(r)|e^{i\varphi(r)}$$


$$v = (\hbar/m)\nabla\varphi \quad \text{implies irrotationality,} \quad \nabla \times v=0.$$



$$L = I\omega = 0$$

In supersolids, the density modulation reduces the superfluid behavior:

$$I = (1 - f_s) I_c$$



Is solid helium a supersolid?



PHYSICAL REVIEW A

VOLUME 2, NUMBER 1

JULY 1970


Speculations on Bose-Einstein Condensation and Quantum Crystals*

G. V. Chester

Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, New York 14850


(Received 13 May 1969)

It is shown, by almost rigorous arguments, that there exist many-body states of a system of interacting bosons which exhibit both crystalline order and Bose-Einstein condensation into the zero-momentum eigenstate of the single-particle density matrix. The implications of this result are discussed in relation to theories of superfluidity and the nature of quantum crystals.




The large zero-point motion in a solid of light bosonic atoms allows the atoms to exchange their positions.

Bose-Einstein condensate of "defectons"



Reviews: Balibar, Nature 464, 176 (2010); Chan, Hallock, Reatto, J. Low. Temp. Phys. 172, 317 (2013).

Experiments on solid helium



Resonant period of the torsion oscillator:



$$\tau = 2\pi\sqrt{I/K}$$

τ : oscillation period
 I : moment of inertia
 K : elastic constant


E. Kim and M. H. W. Chan, Probable observation of a supersolid helium phase, Nature 427, 225 (2004)

$$I = (1 - f_s) I_c$$

large $f_s \sim 10^{-2}$, in disagreement with Leggett predictions (10^{-4})



Experiments on solid helium




J. Day, J. Beamish, Low-temperature shear modulus changes in solid ⁴He and connection to supersolidity, Nature 450, 853 (2007).

D. Y. Kim and M. H. W. Chan, Absence of supersolidity in solid helium in porous Vycor glass, Phys. Rev. Lett. 109, 155301 (2012).


Problem: the change of period might be explained with a change of the elastic constant of He.

$$\tau = 2\pi\sqrt{I/K}$$

Dislocations change state when lowering the temperature, and the crystal stiffens (K increases).





From single particle to cluster supersolids



Helium has a hard-core interaction potential.
There is space only for one particle.
Particles exchange relies only on kinetic energy.

Soft-core interaction potentials allow supersolids with more than one particle per site.




- finite energy cost for multiple occupation of a site
- repulsion between sites, lattice constant $d \approx R_c$
- phase transitions by tuning the interaction strength U


Y. Pomeau and S. Rica, dynamics of a model of a supersolid, Phys. Rev. Lett. 72, 2426 (1994)

Searching cluster supersolids

Weak dressing of Rydberg levels in a Bose-Einstein condensate.



Infinite range interactions for a BEC in optical cavities.



J. Léonard et al., Monitoring and manipulating Higgs and Goldstone modes in a supersolid quantum gas, *Science* 358, 1415 (2017).

Similarly for BECs with light-induced spin-orbit coupling.

J. R. Li et al., A stripe phase with supersolid properties in spin-orbit-coupled Bose-Einstein condensates, *Nature* 543 (2017)

N. Helkel et al., 3D roton excitations and supersolid formation in Rydberg-excited Bose-Einstein condensates, *Phys. Rev. Lett.* 104, 195302 (2010).

Supersolid phase in Bose-Einstein condensates of magnetic atoms

PHYSICAL REVIEW LETTERS



Highlights Recent Accepted Collections Authors Referees Search Press About

Featured in Physics Editors' Suggestion

Observation of a Dipolar Quantum Gas with Metastable Supersolid Properties



L. Tanzi, E. Lucioni, F. Famà, J. Catani, A. Fioretti, C. Gabbanini, R. N. Bisset, L. Santos, and G. Modugno
Phys. Rev. Lett. **122**, 130405 – Published 3 April 2019

Physics See Viewpoint: [Dipolar Quantum Gases go Supersolid](#)




Interactions in dipolar BECs

Interaction between two polarized magnetic dipoles:



Attractive
 $\theta = 0$



Repulsive
 $\theta = \pi/2$

$$U(r) = \frac{4\pi\hbar^2 a_s}{m} \delta(r) + \frac{\mu_0\mu^2}{4\pi} \frac{1 - 3\cos^2\theta}{r^3}$$



van der Waals

dipole-dipole

a_s contact scattering length
Tuneable (Feshbach resonance)

$a_{dd} = \frac{m\mu_0\mu^2}{12\pi\hbar^2}$ dipolar length
Fixed.

Excitation spectrum of dipolar BECs




L. Santos, G. V. Shlyapnikov, and M. Lewenstein, Roton-Maxon spectrum and stability of trapped dipolar Bose-Einstein condensates, Phys. Rev. Lett. 90, 250403 (2003).


Phase diagram

dipolar int. / contact int.
→


BEC: superfluid, follows hydrodynamic equations as an ideal liquid



Droplet crystal: normal solid of self-bound droplets (individually superfluid)





Supersolid?




Experimental methods for dipolar BECs

Typical BECs: $N= 5 \times 10^4$, $T < 50$ nK



Anisotropic optical trap




Dysprosium:
 $\mu = 10 \mu_B$



Melting point:
1680 K


Phase diagram

dipolar int. / contact int.




BEC (superfluid)

Droplet crystal (solid)




Supersolid?



Innsbruck: L. Chomaz et al., Nat. Phys. 14, 442 (2018); D. Petter et al. Phys. Rev. Lett. 122, 183401 (2019).

Stuttgart: H. Kadau et al., Nature 530, 194 (2016); I. Ferrier-Barbut et al., Phys. Rev. Lett. 116, 215301 (2016); M. Wenzel et al., Phys. Rev. A 96, 053630 (2017).

Observation of a coherent density-modulated regime




Stable BEC

Coherent regime (supersolid)


Incoherent regime (droplet crystal)

Experiment: momentum distribution (double-slit interference)

Theory: spatial distribution





3-4 lattice sites,
10⁴ atoms per site



Finite lifetime

Atomic gases decay via formation of molecules.



Transient supersolid properties

PHYSICAL REVIEW LETTERS



Highlights Recent Accepted Collections Authors Referees Search Press About

Featured in Physics Editors' Suggestion

Observation of a Dipolar Quantum Gas with Metastable Supersolid Properties

L. Tanzi, E. Lucioni, F. Famà, J. Catani, A. Fioretti, C. Gabbanini, R. N. Bisset, L. Santos, and G. Modugno
 Phys. Rev. Lett. **122**, 130405 – Published 3 April 2019

Physics See Viewpoint: [Dipolar Quantum Gases go Supersolid](#)




Innsbruck: L. Chomaz et al., Long-lived and transient supersolid behaviors in dipolar quantum gases, Phys. Rev. X 9, 021012 (2019).

Stuttgart: F. Böttcher et al, Transient supersolid properties in an array of dipolar quantum droplets, Phys. Rev. X 9, 011051 (2019).


Spontaneous breaking of two symmetries

Symmetry breaking and Goldstone modes


Modern treatment of sound modes: a gapless Goldstone mode arises each time that an underlying continuous symmetry is spontaneously broken.



Superfluid: gauge symmetry
(phase invariance)




Supersolid: gauge symmetry
and translational symmetry




Solid: translational symmetry

S. Sacconi, S. Moroni and M. Boninsegni, Excitation spectrum of a supersolid, Phys. Rev. Lett. 108, 175301 (2012).


Superfluid regime



Quench of a_s through Feshbach resonances excites the axial breathing mode.




Observable: second moment along x



Frequencies can be measured with relatively high accuracy


$$\omega \sim \sqrt{5/2} \omega_x$$

Symmetry breaking in a supersolid



Two different frequencies for the interference period and interference depth.

Evidence of the lattice compressibility!




lattice mode

superfluid mode

Symmetry breaking in a supersolid

Two quantum phase transitions tuning a single parameter!



L. Tanzi, S. Roccuzzo et al., Supersolid symmetry breaking from compressional oscillations in a dipolar quantum gas, Nature 574, 382 (2019).

Superfluidity from non-classical rotational inertia

Can a Solid Be “Superfluid”?

A. J. Leggett

School of Mathematical and Physical Sciences, University of Sussex, Falmer, Brighton, Sussex, England



(Received 15 September 1970)

Density modulation \Rightarrow local phase variation,
kinetic energy contribution to the free energy:

$$\frac{\hbar^2}{2m} \int (\nabla\varphi)^2 \rho(\vec{r}) d\vec{r} \quad \text{with} \quad \varphi(x) = \int_0^x dx' / \bar{\rho}(x')$$


Reduced superfluid behavior under rotation: $I = (1 - f_s) I_c$

Superfluid fraction: $f_s \simeq \left(\int dx / \bar{\rho}(x) \right)^{-1}$



Rotation of supersolids

Intuitive explanation




A: normal solid

B: supersolid

C: homogeneous superfluid

Caveat: Leggett's approach is one-dimensional and does not consider the superfluidity of individual lattice sites ...

Rotations in an anisotropic trap: scissors mode



Trapped superfluids:
small-angle rotational
oscillation in the
harmonic trap.

In nuclear physics:
relative oscillation of
neutrons and protons

Moment of inertia from oscillation frequency,
similar to torsion oscillators:

$$I = I_c \alpha \beta \frac{(\omega_x^2 + \omega_y^2)}{\omega_{sc}^2}$$

Geometrical
factors:

$$\alpha = (\omega_y^2 - \omega_x^2) / (\omega_x^2 + \omega_y^2)$$

$$\beta = \langle y^2 - x^2 \rangle / \langle y^2 + x^2 \rangle$$

Scissors Mode and Superfluidity of a Trapped Bose-Einstein Condensed Gas

D. Guéry-Odelin and S. Stringari

Dipartimento di Fisica, Università di Trento, and Istituto Nazionale per la Fisica della Materia, I-38050 Povo, Italy
(Received 16 July 1999)

We investigate the oscillation of a dilute atomic gas generated by a sudden rotation of the confining trap (scissors mode). This oscillation reveals the effects of superfluidity exhibited by a Bose-Einstein condensate. The scissors mode is also investigated in a classical gas above T_c in various collisional regimes. The crucial difference with respect to the superfluid case arises from the occurrence of low frequency components, which are responsible for the rigid value of the moment of inertia. Different experimental procedures to excite the scissors mode are discussed.


Observation of the Scissors Mode and Evidence for Superfluidity of a Trapped Bose-Einstein Condensed Gas

O. M. Maragò, S. A. Hopkins, J. Arlt, E. Hodby, G. Hechenblaikner, and C. J. Foot

Clarendon Laboratory, Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom
(Received 18 October 1999)

We report the observation of the scissors mode of a Bose-Einstein condensed gas of ^{87}Rb atoms in a magnetic trap, which gives direct evidence of superfluidity in this system. The scissors mode of oscillation is excited by a sudden rotation of the anisotropic trapping potential. For a gas above T_c (normal fluid) we detect the occurrence of oscillations at two frequencies, with the lower frequency corresponding to the rigid body value of the moment of inertia. Well below T_c the condensate oscillates at a single frequency, without damping, as expected for a superfluid.

Rotations in an anisotropic trap: scissors mode



Trapped superfluids :
small-angle rotational
oscillation in the
harmonic trap.

In nuclear physics:
relative oscillation of
neutrons and protons

Moment of inertia from oscillation frequency,
similar to torsion oscillators:

$$I = I_c \alpha \beta \frac{(\omega_x^2 + \omega_y^2)}{\omega_{sc}^2}$$

Geometrical
factors:

$$\alpha = (\omega_y^2 - \omega_x^2) / (\omega_x^2 + \omega_y^2)$$

$$\beta = \langle y^2 - x^2 \rangle / \langle y^2 + x^2 \rangle$$

For a supersolid:


- Same results for the moment of inertia
- Modified definition of the superfluid fraction, due to the deviation from cylindrical symmetry:

$$I = (1 - f_s) I_c + f_s \beta^2 I_c$$




A.L. Fetter, Vortex nucleation in deformed rotating cylinders.
J. Low Temp. Phys. 16, 533 (1974).

Experimental techniques


The scissors mode is excited by a quench of the strength of the optical traps.



Single mode oscillation in both BEC and supersolid regimes, with different frequencies




Scissors frequency and moment of inertia




L. Tanzi et al. Evidence of superfluidity in a dipolar supersolid from non-classical rotational inertia, arXiv:1912.01910, submitted to Science.
 S. Roccuzzo, A Gallelli, S. Stringari, A. Recati, Rotating a supersolid dipolar gas, Phys. Rev. Lett. 124, 045702 (2020).

Superfluid fraction




Superfluid fraction: $I = (1 - f_s)I_c + f_s \beta^2 I_c \Rightarrow f_s = \frac{1 - \alpha \beta (\omega_x^2 + \omega_y^2) / \omega_{sc}^2}{1 - \beta^2}$

Superfluid fraction: Leggett mechanism




Scissors velocity field (rotating frame)

Leggett formula: $f_s \leq \left(\int dx / \bar{\rho}(x) \right)^{-1}$




Superfluid fraction: Leggett mechanism




Qualitative agreement with Leggett's 1D prediction: large overlap of neighbouring droplets.
 For helium there was a disagreement by 2-3 orders of magnitude.
 Disagreement justified by superfluidity of droplets ...

Rotation of superfluids


Helium-4 (1967)




Helium-3 (1982)




related Meissner effect in superconductors



BECs (2000)



Dipolar supersolid (2020)



Conclusions and outlook


A quantum gas allows studying a fundamental phase of matter intermediate between a liquid (superfluid) and a solid.

Our minimal supersolid is fully controllable and allows testing fundamental properties developed conceptually more than 50 years ago, using methods from various fields of physics.

An unknown quantum material. An extreme form of quantum simulations: testing the properties of future materials...

Future goals:

- Realize larger, 2D supersolids
- Measure a sub-unity superfluid fraction.
- Study the formation of vortices (angular momentum per particle $L = \hbar f_s$).
- Characterize the solid properties of the supersolid (rigidity, plasticity).
- Can a supersolid pass through a capillar without friction?
- Josephson effect without a barrier, ...
- Supersolids in fermionic matter? (pair-density waves in superconductors)



The team – CNR-INO, sede di Pisa and LENS, Università di Firenze



Former members: Francesca Famà, Julian Maloberti, Eleonora Lucioni, Jacopo Catani

Theory by: Russell Bisset, Luis Santos (Hannover); Alessio Recati, Sandro Stringari (Trento); Michele Modugno (Bilbao); Luca Pezzè, Augusto Smerzi (Firenze); Maria Luisa Chiofalo (Pisa), Adriano Angelone (Trieste) ...

<http://quantumgases.lens.unifi.it/exp/dy>



INO-CNR
ISTITUTO
NAZIONALE DI
OTTICA